

Land Surface: Part II

Introduction to warm processes (Vegetation & carbon)

Souhail Boussetta
& the land surface team
souhail.boussetta@ecmwf.int
Room 1114

Outlines

- **Vegetation**
 - **Role of vegetation in NWP**
 - **Tiled approach and current data**
 - **Evolution of vegetation parametrization and practical cases**
- **Carbon**
 - **Why are we interested in carbon?**
 - **Parametrization and feedback from the atmosphere**
 - **Comparison with Jarvis approach and interaction with the atmosphere**

Vegetation state affects

Energy/water budgets

Evapotranspiration

Interception evaporation

Surface albedo (net radiation at the surface)

Aerodynamic exchange through surface roughness

Carbon budget

Plant Respiration

photosynthesis

Vegetation role: some recaps

- Energy balance equation

$$(1-a)R_S^\downarrow + \varepsilon_g R_T^\downarrow - \varepsilon_g \sigma T_{sk}^4 + H + \lambda E = G$$

→ Albedo (a) and emissivity (ε) depend on the surface/vegetation condition




Table 3.1

Radiative Properties of Natural Surfaces^a

Surface type	Other specifications	Albedo (a)	Emissivity (ε)
Water	Small zenith angle	0.03–0.10	0.92–0.97
	Large zenith angle	0.10–0.50	0.92–0.97
Snow	Old	0.40–0.70	0.82–0.89
	Fresh	0.45–0.95	0.90–0.99
Ice	Sea	0.30–0.40	0.92–0.97
	Glacier	0.20–0.40	
Bare sand	Dry	0.35–0.45	0.84–0.90
	Wet	0.20–0.30	0.91–0.95
Bare soil	Dry clay	0.20–0.35	0.95
	Moist clay	0.10–0.20	0.97
	Wet fallow field	0.05–0.07	
Paved	Concrete	0.17–0.27	0.71–0.88
	Black gravel road	0.05–0.10	0.88–0.95
Grass	Long (1 m)	0.16–0.26	0.90–0.95
	Short (0.02 m)		
Agricultural	Wheat, rice, etc.	0.10–0.25	0.90–0.99
	Orchards	0.15–0.20	0.90–0.95
Forests	Deciduous	0.10–0.20	0.97–0.98
	Coniferous	0.05–0.15	0.97–0.99

^a Compiled from Sellers (1965), Kondratyev (1969), and Oke (1978).


Arya, 1988

Vegetation role: some recaps

- Energy balance equation

$$(1-a)R_S^\downarrow + \varepsilon_g R_T^\downarrow - \varepsilon_g \sigma T_{sk}^4 + \textcolor{red}{H} + \lambda E = G$$

→ Sensible heat (H) is also related to vegetation through its relative partition with LE and the aerodynamic exchange specific to surface/vegetation type



Sensible heat flux

$$H = \rho C_h u_L (C_p T_L + gz - C_p T_{sk})$$

$$C_h = f(Ri_B, z_{oh}, z_{om})$$

z_{oh}, z_{om} Roughness length for heat and momentum
Dependent on surface/vegetation type

Vegetation role: some recaps

- **Energy balance equation**


$$(1-a)R_S^\downarrow + \varepsilon_g R_T^\downarrow - \varepsilon_g \sigma T_{sk}^4 + H + \lambda E = G$$

→ Latent heat (LE) is related to vegetation through:


Evapotranspiration and momentum exchange

Interception evaporation= f(Interception reservoir) → f(LAI))

Wet vegetation



$$\left\{ \begin{array}{l} E = \frac{\rho_a}{r_c + r_a} [q_a - q_{sat}(T_{sk})] \\ r_c = \frac{r_{s,\min}}{LAI} \cdot f_1(R_S^\downarrow) f_2(\bar{\theta}) f_3(D_a) \\ r_a = \frac{1}{C_h u_L}, C_h = f(Ri_B, z_{oh}, z_{om}) \end{array} \right.$$



Vegetation role: some recaps

- **Water balance equation**

$$\partial W / \partial t = P - E - R_o - I - D$$

$\partial W / \partial t$ = change in water storage

P = precipitation

E = evapotranspiration

R_o = runoff

I = infiltration

D = lateral diffusion


Evaporation from:

Bare soil

Interception layer

Root transpiration


Infiltration also depend on through fall amount



8 Vegetation heterogeneity

- Land surface is heterogeneous blend of vegetation at many scales
 - forest/cropland/urban area
 - within forest: different trees/moss/understories
- Most LSMs use set of parallel “plant functional types” (PFTs) with specific properties
 - gridbox mean or tiled
 - Some ecological models treat species competition and dynamics within PFTs
- Properties of PFTs
 - LAI
 - rooting depth
 - roughness
 - albedo
 - emission/absorption of organic compounds

CHTESSEL :a tiles approach




F

Land/vegetation	Sea and ice
High vegetation	Open sea / unfrozen lakes
Low vegetation	Sea ice / frozen lakes
High vegetation with snow	
Snow on low vegetation	
Bare ground	
Interception layer	


CHTESSEL geographic characteristics

Fields	ERA15	TESSEL	CHTESSEL
Vegetation	Fraction	Fraction of low Fraction of high	Fraction of low Fraction of high
Vegetation type	Global constant (grass)	Dominant low type Dominant high type	Dominant low type Dominant high type
Albedo	Annual	Monthly	Monthly
LAI	Global constants	Annual, Dependent on vegetation type	Monthly
r_{smin}			
Root depth	1 m	Annual, Dependent on vegetation type	Annual, Dependent on vegetation type
Root profile	Global constant	Annual, Dependent on vegetation type	


High vegetation fraction




Low vegetation fraction



High vegetation Types



Low vegetation Types



Aggregated from GLCC 1km

Vegetation types dependent parameters


Index	Vegetation type	H/L	$r_{s,\min}$ (sm^{-1})	L_{41} ($\text{m}^2 \text{m}^{-2}$)	c_{veg}	g_D (hPa^{-1})	a_r	b_r			
1	Crops, mixed farming	L	180	3	0.90	0	5.558	2.614			
2	Short grass	L	110	2	0.85	0	10.739	2.608			
3	Evergreen needleleaf trees	H	500	5	0.90	0.03	6.706	2.175			
4	Deciduous needleleaf trees	H	500	5	0.90	0.03	7.066	1.953			
5	Deciduous broadleaf trees	H	175	5	0.90	0.03	5.990	1.955			
6	Evergreen broadleaf trees	H	240	6	0.99	0.03	7.344	1.303			
7	Tall grass	L	100	2	0.70	0	8.235	1.627			
8	Desert	-	250								
9	Tundra	L	80								
10	Irrigated crops	L	180	Index	Vegetation type	H/L	$r_{s,\min}$ (sm^{-1})	g_D (hPa^{-1})			
11	Semidesert	L	150	1	Crops, mixed farming	L	100	0.90	0	5.558	2.614
12	Ice caps and glaciers	-	-	2	Short grass	L	100	0.85	0	10.739	2.608
13	Bogs and marshes	L	240	3	Evergreen needleleaf trees	H	250	0.90	0.03	6.706	2.175
14	Inland water	-	-	4	Deciduous needleleaf trees	H	250	0.90	0.03	7.066	1.953
15	Ocean	-	-	5	Deciduous broadleaf trees	H	175	0.90	0.03	5.990	1.955
16	Evergreen shrubs	L	225	6	Evergreen broadleaf trees	H	240	0.99	0.03	7.344	1.303
17	Deciduous shrubs	L	225	7	Tall grass	L	100	0.70	0	8.235	1.627
18	Mixed forest/woodland	H	250	8	Desert	-	250	0	0	4.372	0.978
19	Interrupted forest	H	175	9	Tundra	L	80	0.50	0	8.992	8.992
20	Water and land mixtures	L	150	10	Irrigated crops	L	180	0.90	0	5.558	2.614
Era Interim cycle											
11	Semidesert	L	150	11	Semidesert	L	150	0.10	0	4.372	0.978
12	Ice caps and glaciers	-	-	12	Ice caps and glaciers	-	-	-	-	-	-
13	Bogs and marshes	L	240	13	Bogs and marshes	L	240	0.60	0	7.344	1.303
14	Inland water	-	-	14	Inland water	-	-	-	-	-	-
15	Ocean	-	-	15	Ocean	-	-	-	-	-	-
16	Evergreen shrubs	L	225	16	Evergreen shrubs	L	225	0.50	0	6.326	1.567
17	Deciduous shrubs	L	225	17	Deciduous shrubs	L	225	0.50	0	6.326	1.567
18	Mixed forest/woodland	H	250	18	Mixed forest/woodland	H	250	0.90	0.03	4.453	1.631
19	Interrupted forest	H	175	19	Interrupted forest	H	175	0.90	0.03	4.453	1.631
20	Water and land mixtures	L	150	20	Water and land mixtures	L	150	0.60	0	-	-

Current 45r1 cycle


Index	Vegetation type	H/L	$r_{s,\min}$ (sm^{-1})	c_{veg}	g_D (hPa^{-1})	a_r	b_r
11	Semidesert	L	150	0.10	0	4.372	0.978
12	Ice caps and glaciers	-	-	-	-	-	-
13	Bogs and marshes	L	240	0.60	0	7.344	1.303
14	Inland water	-	-	-	-	-	-
15	Ocean	-	-	-	-	-	-
16	Evergreen shrubs	L	225	0.50	0	6.326	1.567
17	Deciduous shrubs	L	225	0.50	0	6.326	1.567
18	Mixed forest/woodland	H	250	0.90	0.03	4.453	1.631
19	Interrupted forest	H	175	0.90	0.03	4.453	1.631
20	Water and land mixtures	L	150	0.60	0	-	-

**More realistic vegetation dynamic:
Seasonal varying Leaf Area Index**


Seasonal Varying Leaf Area Index



Total LAI [$\text{m}^2 \text{ m}^{-2}$] -July MODIS



Total LAI [$\text{m}^2 \text{ m}^{-2}$] -January MODIS



Obtained by the inversion of a 3D radiative transfer model which compute the LAI and FPAR based on the biome type and an atmospherically corrected surface reflectance thanks to a look-up-table

→derived 8years (2000-2008) climatological time serie

Expected LAI impact on screen level Temperature


For vegetated area the evapotranspiration is parameterized as:

$$E_i = \frac{\rho_a}{r_a + r_c} [q_L - q_{\text{sat}}(T_{\text{sk},i})]$$

Where the canopy resistance r_c is defined following Jarvis(1976) as:


$$r_c = \frac{r_{s,\min}}{LAI} f_1(R_s) f_2(\bar{\theta}) f_3(D_a)$$

Where $r_{s,\min}$ is the minimum stomatal resistance, LAI is the leaf area index and f_1 , f_2 , f_3 are respectively function of the downward shortwave radiation R_s , soil moisture θ and vapour deficit D_a



Analysis experiment: fc experiment validation

Sensitivity





T 2m

Setup: T255

14/02/2008 - 1/09/2008

Seasonal LAI vs fixed LAI




Zonal mean impact

The MODIS LAI introduces a consistent warming seen in FC36h (12UTC) due to reduction of LAI in spring, (increasing vegetation resistance to ET).


This has beneficial impact on near surface temperature forecast (green being positive impact in reducing t2m bias by ~0.5degree)

**More and more realistic vegetation dynamic:
Assimilation of Near Real Time LAI/Albedo**



- NRT analysed LAI is able to fairly detect/monitor anomalous year
- The analysed LAI and albedo signal can be covariant mainly during wet year.

Sensible Heat flux




Clim




NRT_LAI - Clim




NRT_ALB_LAI - Clim



NRT_ALB - Clim



2m temperature sensitivity in coupled run



**Even more realistic vegetation dynamic:
Variable vegetation cover**



Februar

May

July

October

y



**Bare-ground/snow cover
(1- Vegetation fraction)**

→ vegetation cover variation based on satellite observation of Leaf Area Index according to a modified Beer-Lambert law with clumping


$$C_{veg} = 1 - e^{-0.5\omega LAI}$$

February



May

July


October



LAI



(Vegetation cover difference)




Green → more vegetation

→ Physically-based seasonal variability of the vegetation cover


Impact in weather forecast mode

2m temperature [$^{\circ}$ C] NUMBERS: FC-OBS errors [K]
 FC:2015-03-13 12:00:00 STEP 72 VT: 2015-03-16 12:00:00
 N=2768 BIAS= -0.7K STDEV= 2.5K MAE= 2.0K
 errors for [north=-75.00, west=-12.50, south=-35.00, east=42.50]



Cold bias on 2m Temperature
4K on average



2m specific humidity [g/kg] NUMBERS: 10*(FC-OBS)/OBS norm.errors [10s of %]
 FC:2015-03-13 12:00:00 STEP 72 VT: 2015-03-16 12:00:00
 N=2436 BIAS= 8.4% STDEV= 24.5% MAE= 16.6%
 errors for [north=-75.00, west=-12.50, south=-35.00, east=42.50]



Moist bias on 2m specific
humidity 1g/kg on average



Weather forecasts sensitivity

→ Check the T 2m and RH on short term forecast fc+72 valid 12 UTC, March 2015



*Sensitivity = CVEG - CTL ,
if >0 => Warming / adding moisture
if <0 => Cooling / removing moisture*

Weather forecasts impact




Impact = $|CTL - analysis| - |CVEG - analysis|$,


if $>0 \Rightarrow$ relative error reduction from the analysis (**positive impact**)

if $<0 \Rightarrow$ relative error increase from the analysis (**negative impact**)

Behind the scene: change in the forest albedo



Forecast Albedo for CTL




CVEG albedo – CTL albedo


→ Change in the vegetation cover is linked with a change in the forest albedo in presence of snow (in this case)

Introducing Land Carbon parametrisation

Why increasing complexity?



Global Carbon Project 2011; Updated from Le Quéré et al. 2009, Nature G; Canadell et al. 2007, PNAS




The stomatal aperture controls the ratio:
Photosynthesis/Transpiration
 according to the environment conditions
 Light, temperature, air humidity,
 soil moisture, atmospheric [CO₂]

CO₂ and water vapour share the same pathway

The land surface natural contribution to the global carbon budget is highly uncertain

- A better representation of the vegetation processes
- And also attempt to reduce uncertainties from the global carbon budget

Land carbon/photosynthesis-based canopy resistance parameterisation



- $A_n = \rho f(\text{soil m}) \Delta\text{CO}_2 / r_c$
- r_c back-calculated from
 - Empirical soil moisture dependence
 - CO₂-gradient ΔCO_2 is also $f(q_{sat} - q)$
 - Net photosynthetic rate A_n
 - $A_{n,\max}$
 - Photosynthetic active Radiation (PAR)
 - temperature
 - [CO₂]


CTESSEL combines HTESSEL (Balsamo et al. 2009) with the A-gs model used within the ISBA-Ags (Calvet et al. 1998) and developed by Jacobs et al. (1996);

- Account for the effect of CO₂ concentration and the interactions between all environment factors on the stomatal aperture.
- Replaces the Jarvis-type stomata conductance by a photosynthesis dependant-type stomata conductance (Jacobs et al. 1996)
- The model can account for the vegetation response to the radiation at the surface, temperature, soil moisture stress
- Vegetation Assimilation of CO₂ can be used to drive a vegetation growth module to simulate LAI
- The Ecosystem Respiration is parameterized as a function of soil temperature, soil moisture and biome type via a reference respiration parameter


Soil Respiration improvement for winter season

$$NEE = A_n - R_{soil}$$

CETESSEL fi-hyy 2006 RMSE= 1.040 BIAS= -0.512 CORR= 0.945 N= 36



CCTESSEL_dev fluxnet.fi-hyy 2006 RMSE= 0.620 BIAS= -0.221 CORR= 0.934 N= 36



Example of NEE (micro moles /m²/s) predicted over the site Fi-Hyy taking the cold process into account (right) and previous simulation (left) by CTESSEL (black line) and observed (red dots)


Feedback from the atmosphere can contribute to improve the physical understanding and adjust the contribution from the surface

Soil Respiration and winter improvement


$$R_{soil} = R_0 Q_{10}^{(0.1(T_{soil}-25))} f_{sm}$$

➡


$$R_{soil} = R_0 e^{-\alpha \cdot Z_{snow}} Q_{10}^{(0.1(T_{soil}-25))} f_{sm}$$



Including a snow attenuation effect on the soil CO₂ emission





Q10 dependance on Temeprature regime





Including a temeperature dependancy on the Q10 parameter (McGuire et al., 1992)

Jarvis Vs photosynthesis-based evapo-transpiration (offline run)



Surface sensible heat flux (W/m²) compared with flux-tower observations over Fr-LBr for HTESSEL (left panel) and CTESSEL (right panel)




Surface latent heat flux (W/m²) compared with flux-tower observations over Fr-LBr for HTESSEL (left panel) and CTESSEL (right panel).


- CTESSEL improves the LE/H simulations (Photosynthesis-based vs Jarvis approach).

LE/H: When “good” is not enough? (Interaction with the atmosphere)

2m T Error differences from the CTL




2m Rh Error differences from the CTL



Having better LE/H heat flux from the surface does not always lead to a better atmospheric prediction → interaction with other processes and compensating errors?

Near Real Time CO₂ concentration modelled in MACC-II

MACC column-averaged dry-air mole fraction of CO₂ [ppm]
September 2013



- CHTESSEL fluxes used in MACC-II (CAMS) to forecast CO₂ atmospheric concentrations (16 km global simulation)
- Green colours highlight effects of photosynthetic uptake by vegetation
- Diurnal cycle (fluxes driven) and Synoptic variability (Weather driven) are crucial elements for simulating the CO₂ of the Earth system.



Agusti-Panareda et al. (2014, ACP), Boussetta et al. (2013 JGR)

Additional thoughts

-  Taking into account realistic vegetation dynamics is important for accurate representation of surface fluxes and eventually better atmospheric predictability.
-  Carbon, Hydrology and Energy cycles are tightly coupled and an integrated treatment of these processes is a challenge to achieve the necessary accuracy in simulating Net Ecosystem Exchange (CO₂ flux) in global models (and as a component of the global carbon budget).
-  Enhanced connections between albedo, LAI (and roughness) in Earth System Models (ESMs) will most likely increase the sensitivity to vegetation dynamics, and with increased surface related satellite observation products there is potential for further improvements of NWP systems linked with land surface. (better initialisation/ better process description/ possibility to better tune non-observable model parameters)
-  With increased resolution ESMs will have to take into account additional layer of physical complexity such as
 -  vegetation interaction with snow/frozen soil,
 -  better vegetation dynamics
 -  surface- atmosphere coupling and the link with satellite LST,
 -  CO₂/evapo-transpiration coupled processes and satellite fluorescence observation

Thank you

souhail.boussetta@ecmwf.int
Room 1114